

Immunocompromised criteria for patients aged 5-11 years.

Individuals with primary or acquired immunodeficiency states at the time of vaccination due to conditions including:

- acute and chronic leukaemias, and clinically aggressive lymphomas (including Hodgkin's lymphoma) who were under treatment or within 12 months of achieving cure at the time of vaccination
- individuals under follow up for a chronic lymphoproliferative disorders including haematological malignancies
- children with immunosuppression due to HIV/AIDS (children with a current CD4 count of <500 cells/ μ l in those aged 5 years and <200 cells/ μ l in those aged 6-11 years)
- Primary or acquired cellular and combined immune deficiencies – those with lymphopaenia (<1,000 lymphocytes/ μ) or with a functional lymphocyte disorder
- those who had received an allogeneic (cells from a donor) or an autologous (using their own cells) stem cell transplant in the 24 months before vaccination
- those who had received a stem cell transplant more than 24 months before vaccination but had ongoing immunosuppression or graft versus host disease (GVHD)
- persistent agammaglobulinaemia (IgG < 3g/L) due to primary immunodeficiency (e.g. common variable immunodeficiency) or secondary to disease / therapy

Individuals on immunosuppressive or immunomodulating therapy at the time of vaccination including:

- those who were receiving immunosuppressive therapy for a solid organ transplant at the time of vaccination
- those who were receiving or had received in the previous 3 months targeted therapy for autoimmune disease, such as JAK inhibitors or biologic immune modulators including B-cell targeted therapies (including rituximab but in this case the recipient would be considered immunosuppressed for a 6 month period), T-cell co-stimulation modulators, monoclonal tumour necrosis factor inhibitors (TNFi), soluble TNF receptors, interleukin (IL)-6 receptor inhibitors, IL-17 inhibitors, IL 12/23 inhibitors, IL 23 inhibitors. (Note: this list is not exhaustive)
- those who were receiving or had received immunosuppressive chemotherapy or radiotherapy for any indication in the 6 months before vaccination

Individuals with chronic immune-mediated inflammatory disease who were receiving or had received immunosuppressive therapy prior to vaccination including:

- high dose corticosteroids (equivalent to $\geq 1\text{mg}$ prednisolone per kg per day) for more than 10 days in the month before vaccination
- long term moderate dose corticosteroids (equivalent to $\geq 0.5\text{ mg}$ prednisolone per kg per day for more than 4 weeks) in the 3 months before vaccination
- any dose of non-biological oral immune modulating drugs (with the exception of hydroxychloroquine and sulfasalazine), such as methotrexate, azathioprine, 6-mercaptopurine or mycophenolate in the 3 months before vaccination. (Note: this list is not exhaustive)

Individuals who had received high dose steroids (equivalent to $>2\text{mg}$ prednisolone per kg per day for more than a week) for any reason in the month before vaccination